

# Connes' Integral Formula from 1915

Oberwolfach 2026

---

Eva-Maria Hekkelman

Max Planck Institute for Mathematics, Bonn

January 29, 2026

# The Protagonists

## Szegő Limit Theorem (1915)

Let  $w \in C^1(\mathbb{T})$  be positive, with Fourier coefficients

$$w_n := \frac{1}{2\pi} \int_0^{2\pi} w(\theta) e^{-in\theta} d\theta.$$

Define the Toeplitz matrices

$$T_n := \begin{pmatrix} w_0 & w_1 & w_2 & \cdots & w_n \\ w_{-1} & w_0 & w_1 & \cdots & w_{n-1} \\ w_{-2} & w_{-1} & w_0 & \cdots & w_{n-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ w_{-n} & w_{-n+1} & w_{-n+2} & \cdots & w_0 \end{pmatrix}.$$

Then,

$$\lim_{n \rightarrow \infty} (\det T_n)^{\frac{1}{n}} = \exp \left( \frac{1}{2\pi} \int_0^{2\pi} \log(w(\theta)) d\theta \right).$$

## The Protagonists II

### Connes' Integral Formula (1988)

Let  $(M, g)$  be a  $d$ -dimensional compact orientable Riemannian manifold. Then for any Dixmier trace  $\text{Tr}_\omega$ ,

$$\text{Tr}_\omega(M_f(1 - \Delta_g)^{-\frac{d}{2}}) = C_d \int_M f d\nu_g, \quad f \in C^\infty(M),$$

where  $\Delta_g$  is the Laplace–Beltrami operator,  $\nu_g$  the Riemannian volume form, and  $C_d$  a constant depending on the dimension  $d$ .

# Summary of this talk

1. 1915: Szegő's limit theorem
2. 1979: Widom's argument
3. 1988: Connes' integral formula
4. 2025 (I): Noncommutative Szegő limit theorem
5. 2025 (II): NCG and Quantum Ergodicity

This talk is based on joint work with Edward McDonald (Université Paris-Est Créteil).

## Part 1: 1915

---

# Toeplitz Matrices

Let  $\{e_n\}_{n \in \mathbb{Z}}$  be the standard basis of  $L_2(\mathbb{T})$ . The matrix elements of the multiplication operator  $M_w$  are

$$\langle e_n, M_w e_m \rangle = \frac{1}{2\pi} \int_0^{2\pi} e^{-in\theta} w(\theta) e^{im\theta} d\theta = w_{n-m}.$$

The matrix elements of  $T_n$  are also of the form  $w_{m-n}$ .

# Toeplitz Matrices

Let  $\{e_n\}_{n \in \mathbb{Z}}$  be the standard basis of  $L_2(\mathbb{T})$ . The matrix elements of the multiplication operator  $M_w$  are

$$\langle e_n, M_w e_m \rangle = \frac{1}{2\pi} \int_0^{2\pi} e^{-in\theta} w(\theta) e^{im\theta} d\theta = w_{n-m}.$$

The matrix elements of  $T_n$  are also of the form  $w_{m-n}$ .

We therefore have

$$T_n = P_n M_w P_n,$$

where  $P_n$  is the orthogonal projection in  $L_2(\mathbb{T})$  onto the Fourier modes  $\{e_0, e_1, \dots, e_n\}$ .

## Reworking Szegő's theorem

Szegő's limit theorem gives

$$\lim_{n \rightarrow \infty} (\det T_n)^{\frac{1}{n}} = \exp \left( \frac{1}{2\pi} \int_0^{2\pi} \log(w(\theta)) d\theta \right).$$

# Reworking Szegő's theorem

Szegő's limit theorem gives

$$\lim_{n \rightarrow \infty} (\det T_n)^{\frac{1}{n}} = \exp \left( \frac{1}{2\pi} \int_0^{2\pi} \log(w(\theta)) d\theta \right).$$

We have

$$\log(\det(T)) = \log \left( \prod_{\lambda_j \in \sigma(T)} \lambda_j \right) = \sum_{\lambda_j \in \sigma(T)} \log(\lambda_j) = \text{Tr}(\log(T)),$$

# Reworking Szegő's theorem

Szegő's limit theorem gives

$$\lim_{n \rightarrow \infty} (\det T_n)^{\frac{1}{n}} = \exp \left( \frac{1}{2\pi} \int_0^{2\pi} \log(w(\theta)) d\theta \right).$$

We have

$$\log(\det(T)) = \log \left( \prod_{\lambda_j \in \sigma(T)} \lambda_j \right) = \sum_{\lambda_j \in \sigma(T)} \log(\lambda_j) = \text{Tr}(\log(T)),$$

hence a different way to put Szegő's theorem is

$$\frac{1}{n+1} \text{Tr}(\log(P_n M_w P_n)) \xrightarrow{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log(w(\theta)) d\theta,$$

where  $P_n$  is the orthogonal projection in  $L_2(\mathbb{T})$  onto the Fourier modes  $\{e_0, e_1, \dots, e_n\}$ .

## Szegő's limit theorem v2

In fact, Szegő proved a stronger statement.

### Szegő's limit theorem (1915)

For  $0 < w \in C^1(\mathbb{T})$ ,

$$\frac{1}{n+1} \text{Tr}(f(P_n M_w P_n)) \xrightarrow{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} f(w(\theta)) d\theta, \quad f \in C(\mathbb{R}).$$

# Truncated Spectral Triples

---

Spoiler warning!

# Truncated Spectral Triples

Spoiler warning!

If we have a spectral triple  $(\mathcal{A}, \mathcal{H}, D)$ , lots of noncommutative geometers are interested in truncated triples  $(P\mathcal{A}P, P\mathcal{H}, PD)$ , where e.g.  $P := \chi_I(D)$  a finite-rank spectral projection (e.g. Connes–van Suijlekom, D’Andrea–Lizzi–Martinetti).

# Truncated Spectral Triples

Spoiler warning!

If we have a spectral triple  $(\mathcal{A}, \mathcal{H}, D)$ , lots of noncommutative geometers are interested in truncated triples  $(P\mathcal{A}P, P\mathcal{H}, PD)$ , where e.g.  $P := \chi_I(D)$  a finite-rank spectral projection (e.g. Connes–van Suijlekom, D’Andrea–Lizzi–Martinetti).

Note that  $P\mathcal{A}P$  is no longer an algebra: it is in general not closed under products. Instead, it is an *operator system*.

## The Game

An interesting program in NCG is to determine what information about the triple  $(\mathcal{A}, \mathcal{H}, D)$  can be recovered from the operator system spectral triple  $(P\mathcal{A}P, P\mathcal{H}, PD)$ , in particular as  $P \uparrow 1$ .

## The Game

An interesting program in NCG is to determine what information about the triple  $(\mathcal{A}, \mathcal{H}, D)$  can be recovered from the operator system spectral triple  $(P\mathcal{A}P, P\mathcal{H}, PD)$ , in particular as  $P \uparrow 1$ .

Actively studied open questions:

- Under what conditions does the state space of  $P\mathcal{A}P$  equipped with the Connes distance (analogous to Wasserstein 1 metric) converge as a metric space to the state space of  $\mathcal{A}$  as  $P \uparrow 1$ ?

## The Game

An interesting program in NCG is to determine what information about the triple  $(\mathcal{A}, \mathcal{H}, D)$  can be recovered from the operator system spectral triple  $(P\mathcal{A}P, P\mathcal{H}, PD)$ , in particular as  $P \uparrow 1$ .

Actively studied open questions:

- Under what conditions does the state space of  $P\mathcal{A}P$  equipped with the Connes distance (analogous to Wasserstein 1 metric) converge as a metric space to the state space of  $\mathcal{A}$  as  $P \uparrow 1$ ?
- Can we recover the  $K$ -theory of  $(\mathcal{A}, \mathcal{H}, D)$  from its truncations?

## Szegő's limit theorem v2 again

In this light and in this form, Szegő's limit theorem looks promising.

### Szegő's limit theorem (1915)

For  $0 < w \in C^1(\mathbb{T})$ ,

$$\frac{1}{n+1} \text{Tr}(f(P_n M_w P_n)) \xrightarrow{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} f(w(\theta)) d\theta, \quad f \in C(\mathbb{R}).$$

## Part 2: 1979

---

## Widom's formula

In the 1970s, Szegő's formula received renewed attention, due to the emergence of the field of Quantum Ergodicity. Widom managed to generalise the formula to manifolds.

# Widom's formula

In the 1970s, Szegő's formula received renewed attention, due to the emergence of the field of Quantum Ergodicity. Widom managed to generalise the formula to manifolds.

## Widom's Szegő's limit theorem (1979)

Let  $(M, g)$  be a compact Riemannian manifold,  $w \in C^\infty(M)$  real-valued. Then

$$\frac{\text{Tr}(f(P_\lambda M_w P_\lambda))}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\text{vol}(M)} \int_M f(w(x)) d\nu_g(x), \quad f \in C(\mathbb{R}),$$

where  $P_\lambda = \chi_{[-\lambda, \lambda]}(\Delta_g)$ .

# Widom's formula

In the 1970s, Szegő's formula received renewed attention, due to the emergence of the field of Quantum Ergodicity. Widom managed to generalise the formula to manifolds.

## Widom's Szegő's limit theorem (1979)

Let  $(M, g)$  be a compact Riemannian manifold,  $w \in C^\infty(M)$  real-valued. Then

$$\frac{\text{Tr}(f(P_\lambda M_w P_\lambda))}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\text{vol}(M)} \int_M f(w(x)) d\nu_g(x), \quad f \in C(\mathbb{R}),$$

where  $P_\lambda = \chi_{[-\lambda, \lambda]}(\Delta_g)$ . Furthermore, for  $A \in \Psi^0(M)$  self-adjoint with principal symbol  $\sigma_0(A)$ ,

$$\frac{\text{Tr}(f(P_\lambda A P_\lambda))}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\text{vol}(S^*M)} \int_{S^*M} f(\sigma_0(A)) d\mu, \quad f \in C(\mathbb{R}).$$

# Idea

Widom's proof is quite short.

- The first step is to prove the case where  $f(x) = x$ , i.e.

$$\frac{\text{Tr}(P_\lambda A P_\lambda)}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu,$$

which is in fact the microlocal Weyl law.

# Idea

Widom's proof is quite short.

- The first step is to prove the case where  $f(x) = x$ , i.e.

$$\frac{\text{Tr}(P_\lambda A P_\lambda)}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu,$$

which is in fact the microlocal Weyl law.

- The next step is to do polynomials, by proving that

$$\frac{\text{Tr}((P_\lambda A P_\lambda)^n - P_\lambda A^n P_\lambda)}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} 0.$$

## Microlocal Weyl law

Weyl's law gives for a compact Riemannian manifold  $(M, g)$ ,

$$\mathrm{Tr}(P_\lambda) \sim C_d \mathrm{vol}(M) \lambda^{\frac{d}{2}}, \quad \lambda \rightarrow \infty.$$

## Microlocal Weyl law

Weyl's law gives for a compact Riemannian manifold  $(M, g)$ ,

$$\mathrm{Tr}(P_\lambda) \sim C_d \mathrm{vol}(M) \lambda^{\frac{d}{2}}, \quad \lambda \rightarrow \infty.$$

There exists a *local* version of Weyl's law, which gives for  $f \in C^\infty(M)$ ,

$$\mathrm{Tr}(M_f P_\lambda) = \sum_{n=0}^{N(\lambda)} \langle e_n, M_f e_n \rangle \sim C_d \lambda^{\frac{d}{2}} \int_M f d\nu_g, \quad \lambda \rightarrow \infty.$$

## Microlocal Weyl law

Weyl's law gives for a compact Riemannian manifold  $(M, g)$ ,

$$\mathrm{Tr}(P_\lambda) \sim C_d \mathrm{vol}(M) \lambda^{\frac{d}{2}}, \quad \lambda \rightarrow \infty.$$

There exists a *local* version of Weyl's law, which gives for  $f \in C^\infty(M)$ ,

$$\mathrm{Tr}(M_f P_\lambda) = \sum_{n=0}^{N(\lambda)} \langle e_n, M_f e_n \rangle \sim C_d \lambda^{\frac{d}{2}} \int_M f d\nu_g, \quad \lambda \rightarrow \infty.$$

Or, even, a *microlocal* Weyl law, which states for  $A \in \Psi^0(M)$ ,

$$\mathrm{Tr}(A P_\lambda) = \sum_{n=0}^{N(\lambda)} \langle e_n, A e_n \rangle \sim C_d \lambda^{\frac{d}{2}} \int_{S^* M} \sigma_0(A) d\mu, \quad \lambda \rightarrow \infty.$$

## Part 3: 1988

---

## Dixmier traces

Let  $\mathcal{H}$  be a Hilbert space. An eigenvalue sequence of a compact operator  $A \in K(\mathcal{H})$  is a sequence  $\{\lambda(k, A)\}_{k \in \mathbb{N}}$  of the eigenvalues of  $A$  listed with multiplicity, such that  $\{|\lambda(k, A)|\}_{k \in \mathbb{N}}$  is non-increasing.

The usual operator trace  $\text{Tr}$  can be characterised for trace class operators  $A \in \mathcal{L}_1 \subset K(\mathcal{H})$  as

$$\text{Tr}(A) = \lim_{n \rightarrow \infty} \sum_{k=1}^n \lambda(k, A).$$

## Dixmier traces

Let  $\mathcal{H}$  be a Hilbert space. An eigenvalue sequence of a compact operator  $A \in K(\mathcal{H})$  is a sequence  $\{\lambda(k, A)\}_{k \in \mathbb{N}}$  of the eigenvalues of  $A$  listed with multiplicity, such that  $\{|\lambda(k, A)|\}_{k \in \mathbb{N}}$  is non-increasing.

The usual operator trace  $\text{Tr}$  can be characterised for trace class operators  $A \in \mathcal{L}_1 \subset K(\mathcal{H})$  as

$$\text{Tr}(A) = \lim_{n \rightarrow \infty} \sum_{k=1}^n \lambda(k, A).$$

The Dixmier trace is defined on so-called weak trace class operators  $A \in \mathcal{L}_{1,\infty} \subset K(\mathcal{H})$  by

$$\text{Tr}_\omega(A) = \omega\text{-}\lim_{n \rightarrow \infty} \frac{1}{\log(2+n)} \sum_{k=1}^n \lambda(k, A),$$

where  $\omega \in \ell_\infty(\mathbb{N})^*$  is an extended limit. Note that  $\mathcal{L}_1 \subset \mathcal{L}_{1,\infty}$ , but if  $A \in \mathcal{L}_1$ ,  $\text{Tr}_\omega(A) = 0$ .

# Connes' integral formula

Connes proved the following.

## Connes' Integral Formula

Let  $(M, g)$  be a compact Riemannian manifold,  $f \in C_c^\infty(M)$ . Then for any Dixmier trace  $\text{Tr}_\omega$ ,

$$\text{Tr}_\omega(M_f(1 - \Delta_g)^{-\frac{d}{2}}) = C_d \int_M f d\nu_g.$$

# Connes' integral formula

Connes proved the following.

## Connes' Integral Formula

Let  $(M, g)$  be a compact Riemannian manifold,  $f \in C_c^\infty(M)$ . Then for any Dixmier trace  $\text{Tr}_\omega$ ,

$$\text{Tr}_\omega(M_f(1 - \Delta_g)^{-\frac{d}{2}}) = C_d \int_M f d\nu_g.$$

Or stronger, for  $A \in \Psi_{cl}^0(M)$ ,

$$\text{Tr}_\omega(A(1 - \Delta_g)^{-\frac{d}{2}}) = C_d \int_{S^*M} \sigma_0(A) d\mu.$$

# Connes' integral formula

Connes proved the following.

## Connes' Integral Formula

Let  $(M, g)$  be a compact Riemannian manifold,  $f \in C_c^\infty(M)$ . Then for any Dixmier trace  $\text{Tr}_\omega$ ,

$$\text{Tr}_\omega(M_f(1 - \Delta_g)^{-\frac{d}{2}}) = C_d \int_M f d\nu_g.$$

Or stronger, for  $A \in \Psi_{cl}^0(M)$ ,

$$\text{Tr}_\omega(A(1 - \Delta_g)^{-\frac{d}{2}}) = C_d \int_{S^*M} \sigma_0(A) d\mu.$$

Connes' result is in fact even stronger, as he does not assume a Riemannian structure.

## Part 4: 2025 (I)

---

## Comparison

Now compare the first step in Widom's proof, the microlocal Weyl law

$$\frac{\mathrm{Tr}(P_\lambda A P_\lambda)}{\mathrm{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\mathrm{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu,$$

with Connes' formula

$$\mathrm{Tr}_\omega(A(1 - \Delta)^{-\frac{d}{2}}) = C_d \int_{S^*M} \sigma_0(A) d\mu.$$

# Comparison

Now compare the first step in Widom's proof, the microlocal Weyl law

$$\frac{\text{Tr}(P_\lambda AP_\lambda)}{\text{Tr}(P_\lambda)} \xrightarrow{\lambda \rightarrow \infty} \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu,$$

with Connes' formula

$$\text{Tr}_\omega(A(1 - \Delta)^{-\frac{d}{2}}) = C_d \int_{S^*M} \sigma_0(A) d\mu.$$

## H.-McDonald

Let  $\mathcal{H}$  be a separable Hilbert space,  $A \in B(\mathcal{H})$ ,  $D$  self-adjoint with compact resolvent,  $P_\lambda := \chi_{[-\lambda, \lambda]}(D)$ . If  $D$  satisfies Weyl's law  $\lambda(k, |D|) \sim Ck^{\frac{1}{d}}$ , then for all Dixmier traces  $\text{Tr}_\omega$ ,

$$\frac{\text{Tr}_\omega(A(1 + D^2)^{-\frac{d}{2}})}{\text{Tr}_\omega((1 + D^2)^{-\frac{d}{2}})} = \omega \circ M \left( \frac{\text{Tr}(P_{\lambda_n} AP_{\lambda_n})}{\text{Tr}(P_{\lambda_n})} \right).$$

Here,  $M : \ell_\infty \rightarrow \ell_\infty$  is the logarithmic averaging defined by  $M(x)_n = \frac{1}{\log(n+2)} \sum_{k=0}^n \frac{x_k}{k}$ .

# Truncated Spectral Triples

Our result shows that if  $(\mathcal{A}, \mathcal{H}, D)$  is  $d$ -dimensional and  $D$  satisfies Weyl's law, then

$$P_\lambda A P_\lambda \mapsto \frac{\text{Tr}(P_\lambda A P_\lambda)}{\text{Tr}(P_\lambda)}$$

is a reasonable approximation of the noncommutative integral  $\text{Tr}_\omega(A(1+D^2)^{-\frac{d}{2}})$ .

# Noncommutative Szegő theorem

With Widom's argument, we have a Szegő formula for NCG as well.

## H.-McDonald

Let  $\mathcal{H}$  be a separable Hilbert space,  $A \in B(\mathcal{H})_{sa}$ ,  $D$  self-adjoint with compact resolvent. If  $D$  satisfies Weyl's law  $\lambda(k, |D|) \sim Ck^{\frac{1}{d}}$ , and if  $[D, A]$  extends to a bounded operator, then for all Dixmier traces  $\text{Tr}_\omega$ ,

$$\frac{\text{Tr}_\omega(f(A)(1+D^2)^{-\frac{d}{2}})}{\text{Tr}_\omega((1+D^2)^{-\frac{d}{2}})} = \omega \circ M \left( \frac{\text{Tr}(f(P_{\lambda_n} A P_{\lambda_n}))}{\text{Tr}(P_{\lambda_n})} \right), \quad f \in C(\mathbb{R}).$$

# Noncommutative Szegő theorem

With Widom's argument, we have a Szegő formula for NCG as well.

## H.-McDonald

Let  $\mathcal{H}$  be a separable Hilbert space,  $A \in B(\mathcal{H})_{sa}$ ,  $D$  self-adjoint with compact resolvent. If  $D$  satisfies Weyl's law  $\lambda(k, |D|) \sim Ck^{\frac{1}{d}}$ , and if  $[D, A]$  extends to a bounded operator, then for all Dixmier traces  $\text{Tr}_\omega$ ,

$$\frac{\text{Tr}_\omega(f(A)(1+D^2)^{-\frac{d}{2}})}{\text{Tr}_\omega((1+D^2)^{-\frac{d}{2}})} = \omega \circ M \left( \frac{\text{Tr}(f(P_{\lambda_n}AP_{\lambda_n}))}{\text{Tr}(P_{\lambda_n})} \right), \quad f \in C(\mathbb{R}).$$

In case the spectral triple has *local Weyl laws*, i.e.  $\text{Tr}(aP_\lambda) \sim C(a)\lambda^{\frac{d}{2}}$  for all  $a \in \mathcal{A}$ , we have

$$\frac{\text{Tr}_\omega(f(a)(1+D^2)^{-\frac{d}{2}})}{\text{Tr}_\omega((1+D^2)^{-\frac{d}{2}})} = \lim_{\lambda \rightarrow \infty} \frac{\text{Tr}(f(P_\lambda a P_\lambda))}{\text{Tr}(P_\lambda)}, \quad f \in C(\mathbb{R}).$$

# Noncommutative Szegő theorem

In particular, in this setting we recover Szegő's limit theorem

$$\exp \left( \frac{\mathrm{Tr}_\omega(\log(a)(1+D^2)^{-\frac{d}{2}})}{\mathrm{Tr}_\omega((1+D^2)^{-\frac{d}{2}})} \right) = \lim_{\lambda \rightarrow \infty} (\det(P_\lambda a P_\lambda))^{\mathrm{Tr}(P_\lambda)}, \quad 0 < a \in \mathcal{A}.$$

## Part 5: 2025 (II)

---

# Quantum Ergodicity


The result by Szegő was studied by Widom in the context of Quantum Ergodicity.

# Quantum Ergodicity


The result by Szegő was studied by Widom in the context of Quantum Ergodicity.

This field studies a quantum mechanical analogue of ergodicity. A one-particle system described by an operator  $H$  on  $L_2(M)$  is called quantum ergodic if the high energy states of  $H$  are ‘smeared’ over  $M$ .

# Picture



**Figure 1:** Eigenfunctions of the Laplacian on a rose-shaped domain, quantum ergodicity **unknown**.



**Figure 2:** Eigenfunctions of the Laplacian on the disc, **not** quantum ergodic.

## Quantum Ergodicity

For a compact Riemannian manifold  $M$  and a positive self-adjoint operator  $\Delta$  on  $L_2(M)$  with compact resolvent,  $\Delta$  is said to be quantum ergodic if for every orthonormal basis  $\{e_n\}_{n=0}^\infty$  of  $L_2(M)$  consisting of eigenfunctions of  $\Delta$ , there exists a density one subsequence  $J \subseteq \mathbb{N}$  such that

$$\lim_{J \ni j \rightarrow \infty} \langle e_j, A e_j \rangle_{L_2(M)} \rightarrow \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu, \quad A \in \Psi^0(M),$$

where  $\nu$  is a probability measure on  $M$ . In this context, a density one subsequence means that

$$\frac{\#(J \cap \{0, \dots, n\})}{n+1} \rightarrow 1, \quad n \rightarrow \infty.$$

## QE as a Weyl law

We can interpret Quantum Ergodicity as a stronger microlocal Weyl law. Namely, the QE property

$$\lim_{j \ni j \rightarrow \infty} \langle e_j, A e_j \rangle_{L_2(M)} \rightarrow \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu, \quad A \in \Psi^0(M),$$

is equivalent by the Koopman–von Neumann lemma to

$$\lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N \left| \langle e_n, A e_n \rangle - \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu \right| = 0, \quad A \in \Psi^0(M).$$

## QE as a Weyl law

We can interpret Quantum Ergodicity as a stronger microlocal Weyl law. Namely, the QE property

$$\lim_{J \ni j \rightarrow \infty} \langle e_j, A e_j \rangle_{L_2(M)} \rightarrow \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu, \quad A \in \Psi^0(M),$$

is equivalent by the Koopman–von Neumann lemma to

$$\lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N \left| \langle e_n, A e_n \rangle - \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu \right| = 0, \quad A \in \Psi^0(M).$$

This is now recognisable as a stronger version of the microlocal Weyl law

$$\lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N \langle e_n, A e_n \rangle - \frac{1}{\text{vol}(S^*M)} \int_{S^*M} \sigma_0(A) d\mu = 0, \quad A \in \Psi^0(M).$$

## Geodesic flow

Let  $(M, g)$  be a Riemannian manifold, and denote by  $SM \subseteq TM$  the tangent vectors of length 1. Then we define the geodesic flow

$$G_t : SM \rightarrow SM, \quad t \in \mathbb{R},$$

in the usual way. By duality, we can likewise define  $G_t : S^*M \rightarrow S^*M$ , where  $S^*M \subseteq T^*M$ .

## Geodesic flow

Let  $(M, g)$  be a Riemannian manifold, and denote by  $SM \subseteq TM$  the tangent vectors of length 1. Then we define the geodesic flow

$$G_t : SM \rightarrow SM, \quad t \in \mathbb{R},$$

in the usual way. By duality, we can likewise define  $G_t : S^*M \rightarrow S^*M$ , where  $S^*M \subseteq T^*M$ .

The geodesic flow is said to be *ergodic* if for every measurable function  $f \in L_\infty(S^*M)$  which is fixed by the flow (i.e.  $f \circ G_t = f$  almost everywhere), it must be that  $f$  is constant almost everywhere.

# Fundamental theorem of QE

The fundamental theorem that started the field of Quantum Ergodicity is the following.

**Theorem (Shnirelman 1974, Zelditch 1987, Colin de Verdière 1985)**

Let  $M$  be a compact Riemannian manifold. If the geodesic flow on  $M$  is ergodic, then the Laplace–Beltrami operator  $\Delta_g$  is quantum ergodic.

# Fundamental theorem of QE

The fundamental theorem that started the field of Quantum Ergodicity is the following.

## Theorem (Shnirelman 1974, Zelditch 1987, Colin de Verdière 1985)

Let  $M$  be a compact Riemannian manifold. If the geodesic flow on  $M$  is ergodic, then the Laplace–Beltrami operator  $\Delta_g$  is quantum ergodic.

By now, various extensions of this theorem exist. The common thread is to study geodesic flow, and translate this into asymptotic behaviour of eigenfunctions of an operator.

# Noncommutative geodesic flow

Connes defined geodesic flow on spectral triples in 1996.

## Noncommutative cosphere bundle

Let  $(\mathcal{A}, \mathcal{H}, D)$  be a regular spectral triple. Let  $\sigma_t : B(\mathcal{H}) \rightarrow B(\mathcal{H})$  be defined by  $A \mapsto e^{it|D|} A e^{-it|D|}$ . Then define the  $C^*$ -algebra

$$S^* \mathcal{A} := C^* \left( \bigcup_{t \in \mathbb{R}} \sigma_t(\mathcal{A}) + K(\mathcal{H}) \right) \Big/ K(\mathcal{H}),$$

where  $\sigma_t$  descends to an action of  $\mathbb{R}$  on  $S^* \mathcal{A}$ . If  $(\mathcal{A}, \mathcal{H}, D) \simeq (C^\infty(M), L_2(S), D_S)$ , then  $S^* \mathcal{A} \simeq C(S^* M)$  and  $\sigma_t$  is the geodesic flow.

## NCG ergodicity

Since ergodicity of the geodesic flow is a measure theoretic statement, we need to take one more step.

Since ergodicity of the geodesic flow is a measure theoretic statement, we need to take one more step.

## $L_2$ -cosphere bundle

Let  $(\mathcal{A}, \mathcal{H}, D)$  be a regular spectral triple where  $D$  satisfies Weyl's law. Then

$$\tau(A + K(\mathcal{H})) = \frac{\text{Tr}_\omega(A(1 + D^2)^{-\frac{d}{2}})}{\text{Tr}_\omega((1 + D^2)^{-\frac{d}{2}})}, \quad A + K(\mathcal{H}) \in S^* \mathcal{A},$$

defines a finite positive trace on  $S^* \mathcal{A}$ . Then define  $L_2(S^* \mathcal{A})$  as the Hilbert space of the GNS representation of  $S^* \mathcal{A}$  corresponding to  $\tau$ .

The geodesic flow  $\sigma_t$  on  $S^* \mathcal{A}$  descends to a unitary operator on  $L_2(S^* \mathcal{A})$ .

We can now naively put forward a definition of ergodic geodesic flow for spectral triples. Namely, we say that the geodesic flow  $\sigma_t$  is ergodic on  $(\mathcal{A}, \mathcal{H}, D)$  if the only  $\sigma_t$ -invariant element of  $L_2(S^* \mathcal{A})$  is the identity.

We can now naively put forward a definition of ergodic geodesic flow for spectral triples. Namely, we say that the geodesic flow  $\sigma_t$  is ergodic on  $(\mathcal{A}, \mathcal{H}, D)$  if the only  $\sigma_t$ -invariant element of  $L_2(S^* \mathcal{A})$  is the identity.

### NCG QE (H.-McDonald)

Let  $(\mathcal{A}, \mathcal{H}, D)$  be a  $d$ -summable regular spectral triple where  $D$  satisfies Weyl's law, and with local Weyl laws. If the geodesic flow on  $(\mathcal{A}, \mathcal{H}, D)$  is ergodic, then  $D$  is quantum ergodic. That is, for every basis  $\{e_n\}_{n=0}^\infty$  of  $\mathcal{H}$  consisting of eigenvectors of  $D$ , there exists a density one subset  $J \subseteq \mathbb{N}$  such that

$$\lim_{J \ni j \rightarrow \infty} \langle e_j, ae_j \rangle = \frac{\text{Tr}_\omega(a(1+D^2)^{-\frac{d}{2}})}{\text{Tr}_\omega((1+D^2)^{-\frac{d}{2}})}, \quad a \in \mathcal{A}.$$

# Thanks

Thanks for listening!